



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

FOREIGN STATISTICAL REPORTS.

ARGENTINA—*Buenos Ayres*.—Month of April, 1900. Estimated population, 800,000. Total number of deaths, 690, including diphtheria, 14; enteric fever, 20; measles, 2; scarlet fever, 20; whooping cough, 1; smallpox, 5, and 128 from tuberculosis.

BAHAMAS—*Dunmore Town*.—Two weeks ended June 25, 1900. Estimated population, 1,472. One death. No contagious diseases.

Governors Harbor.—Two weeks ended June 23, 1900. Estimated population, 1,500. No deaths and no contagious diseases.

Green Turtle Cay—Abaco.—Two weeks ended June 21, 1900. Estimated population, 3,900. No deaths and no contagious diseases.

Nassau.—Two weeks ended June 26, 1900. Estimated population, 12,000. Number of deaths not reported. No contagious diseases.

FRANCE—*Nice*.—Month of May, 1900. Estimated population, 108,227. Total number of deaths, 165, including enteric fever, 2; whooping cough, 3; smallpox, 1, and 37 from tuberculosis.

Roubaix.—Month of May, 1900. Estimated population, 127,476. Total number of deaths, 188, including diphtheria, 1; enteric fever, 3; measles, 3, and 1 from whooping cough.

Rouen.—Month of May, 1900. Estimated population, 112,657. Total number of deaths, 239, including diphtheria, 2; enteric fever, 3; whooping cough, 1, and 47 from tuberculosis.

GERMANY—*Dresden*.—Month of April, 1900. Estimated population, 405,500. Total number of deaths, 656, including diphtheria, 5; enteric fever, 2; measles, 3; scarlet fever, 1; whooping cough, 4, and 104 from phthisis pulmonalis.

Hanover.—Month of April, 1900. Estimated population, 249,362. Total number of deaths, 313, including diphtheria, 6; measles, 2; scarlet fever, 5, and 59 from tuberculosis.

Magdeburg.—Month of March, 1900. Estimated population, 229,596. Total number of deaths, 454, including diphtheria, 16; scarlet fever, 9; whooping cough, 6, and 34 from tuberculosis.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 33 great towns in England and Wales during the week ended June 9, 1900, correspond to an annual rate of 16.8 a thousand of the aggregate population, which is estimated at 11,610,296. The highest rate was recorded in Liverpool, viz, 24.8, and the lowest in Birkenhead, viz, 8.0.

Bradford.—Two weeks ended June 2, 1900. Estimated population, 291,535. Total number of deaths, 193, including diphtheria, 2; enteric fever, 1; measles, 7; scarlet fever, 4, and 1 from whooping cough.

London.—One thousand three hundred and ninety-eight deaths were registered during the week, including measles 60; scarlet fever, 11; diphtheria, 24; whooping cough, 37; enteric fever, 7, and diarrhea and dysentery, 5. The deaths from all causes correspond to an annual rate of 15.9 a thousand. In Greater London 1,868 deaths were regis-

tered, corresponding to an annual rate of 14.6 a thousand of the population. In the "outer ring" the deaths included 9 from diphtheria, 27 from measles, and 21 from whooping cough.

Ireland.—The average annual death rate represented by the deaths registered during the week ended June 9, 1900, in the 22 principal town districts of Ireland was 19.4 a thousand of the population, which is estimated at 1,062,188. The lowest rate was recorded in Galway, viz, 3.8, and the highest in Newtownards, viz, 51.0 a thousand. In Dublin and suburbs 161 deaths were registered, including enteric fever, 2, and 2 from diphtheria.

Scotland.—The deaths registered in 8 principal towns during the week ended June 9, 1900, correspond to an annual rate of 18.3 a thousand of the population, which is estimated at 1,606,935. The lowest mortality was recorded in Paisley, viz, 13.3, and the highest in Greenock, viz, 21.5 a thousand. The aggregate number of deaths registered from all causes was 564, including diphtheria, 3; measles, 19; scarlet fever, 6; smallpox, 1, and 17 from whooping cough.

SWITZERLAND.—Reports for the two weeks ended June 2, 1900, from 15 cities and towns having an aggregate estimated population of 690,000, show a total of 499 deaths, including diphtheria, 4; measles, 8; whooping cough, 3; smallpox, 2, and 97 from phthisis pulmonalis.